

HIGH SENSITIVITY DETECTION OF TUMOR CELLS IN PERIPHERAL BLOOD OF CARCINOMA PATIENTS BY A REVERSE TRANSCRIPTION PCR METHOD

Bertha Gutierrez, Veit Zieglschmid, Christiane Hollmann, Winfried Albert, and Oliver Böcher

AdnaGen AG, Langenhagen, Germany
Correspondence to: Dr. Oliver Böcher, Ostpassage 7, D-30853
Langenhagen, Germany. Tel: +49 511 72595050, Fax: +49 511 72594040,
E-mail: ob@adnagen.com

Abstract

Occurrence of tumor cells in the peripheral blood of individuals suffering from cancer may serve as an early indication that the primary tumor has evaded from its tissue of origin. Despite defence mechanisms of the organism individual cancer cells may attach in distant regions and form colonies as an initial step of metastasis formation. Blood-borne distant metastasis is the leading cause of cancer-related deaths.

Hence, early detection of metastatic potential can be estimated by detecting tumor cells in the blood circulation. Sensitivity and specificity are the main objectives of any method applied for this purpose. Most highly sensitive techniques described, however, encounter an increasing problem of specificity due to background signalling from illegitimate transcription.

AdnaGen established a method of tumor cell selection and their specific identification and analysis with sensitivity of at least 2 tumor cells in 5 mL blood. To achieve this, we combined the enrichment of tumor cells using a specifically designed antibody mixture with RT-multiplex PCR techniques for the detection of mRNAs encoding for tumor associated genes. Preliminary results of case studies showed occurrence of tumor cells in blood of carcinoma patients indicating a potentially tumor relapse, in some cases several months prior to an elevation of serum tumor markers.

In conclusion, AdnaGen provide a sensitive and specific method to detect disseminated tumor cells in peripheral blood of testicular, breast and colorectal carcinoma patients. This innovative method is an option for clinicians as a predictive tool with respect to metastasis formation and may result in an appropriate selection of patients for adjuvant therapy.